

DOCKET NO.: FO-3-C1

ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
WASHINGTON, DC 20231

CERTIFICATION UNDER 37 CFR 1.10

Date of Deposit: September 1, 1999 Mailing Label Number: EL344800793US

I hereby certify that this 37 CFR 1.53(b) request and the documents referred to as attached therein are being deposited with the United States Postal Service on the date indicated above in an envelope as "Express Mail Post Office to Addressee" service under 37 CFR 1.10 and addressed to the Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

Susan A. Gordon
Name of Person Mailing Paper

Susan A. Gordon
Signature of Person Mailing Paper

JC598 U.S. PTO
09/387809
09/01/99

Dear Sir:

This is a request for filing a new nonprovisional application under 37 CFR § 1.53(b) which claims priority in prior pending provisional application serial no. 60/098,911, filed September 2, 1998.

INVENTOR(S): Elizabeth A. Hodgkins, 24355 La Homa, Yorba Linda, California 92887

FOR: "METHOD AND COMPOSITION TO PROTECT AN OBLIGATE CARNIVORE FROM A DISEASE OF ABNORMAL CARBOHYDRATE METABOLISM"

ENCLOSED ARE:

- ☒ 55 PAGES OF SPECIFICATION, CLAIMS AND ABSTRACT
- ☐ _____ SHEETS OF DRAWING (☐ FORMAL OR ☐ INFORMAL)
- ☐ DECLARATION
- ☐ POWER OF ATTORNEY
- ☐ ASSIGNMENT OF THE INVENTION TO: [Heska Corporation] (Under separate cover letter)
- ☐ SMALL ENTITY STATEMENT
- ☐ IDENTICAL PAPER AND COMPUTER READABLE COPIES OF APPLICATION SEQUENCE LISTING. APPLICANT HEREBY ASSERTS PURSUANT TO 37 CFR §1.821(f) THAT THE CONTENT OF THE PAPER AND COMPUTER READABLE COPIES OF SEQ ID NO: __ THROUGH SEQ ID NO: __ SUBMITTED HERewith ARE IDENTICAL.
- ☐ PRELIMINARY AMENDMENT
- ☐ INFORMATION DISCLOSURE STATEMENT
- ☐ PTO 1449 FORM
- ☐ FOREIGN PRIORITY BENEFITS ARE CLAIMED UNDER 35 USC §119 OF _____ (Country) PATENT APPLICATION SERIAL NO _____, FILED _____.
- ☐ CERTIFIED COPY OF A _____ APPLICATION
- ☒ FILING FEE IS NOT ENCLOSED AT THIS TIME.
- ☐ ENCLOSED IS A CHECK IN THE AMOUNT OF \$ _____ TO COVER THE FILING FEE. SEE FOLLOWING PAGE FOR CALCULATION OF FILING FEE.
- ☐ PLEASE CHARGE DEPOSIT ACCOUNT NO 081930 IN THE AMOUNT OF \$ _____. A DUPLICATE COPY OF THIS SHEET IS ENCLOSED.
- ☐ THE COMMISSIONER IS HEREBY AUTHORIZED TO CHARGE PAYMENT OF THE FOLLOWING FEES ASSOCIATED WITH THIS COMMUNICATION OR CREDIT ANY OVERPAYMENT TO DEPOSIT ACCOUNT NO. 081930. A DUPLICATE COPY OF THIS SHEET IS ENCLOSED.
 - ☐ ANY ADDITIONAL FILING FEES REQUIRED UNDER 37 C.F.R. 1.16.
 - ☐ ANY PATENT APPLICATION PROCESSING FEES UNDER 37 C.F.R. 1.17.

JC604 U.S. PTO
09/01/99

09/01/99 10:00:00

☐ THE COMMISSIONER IS HEREBY AUTHORIZED TO CHARGE PAYMENT OF THE FOLLOWING FEES DURING THE PENDENCY OF THIS APPLICATION OR CREDIT ANY OVERPAYMENT TO DEPOSIT ACCOUNT NO. 081930. A DUPLICATE COPY OF THIS SHEET IS ENCLOSED.

- ☐ ANY PATENT APPLICATION PROCESSING FEES UNDER 37 C.F.R. 1.17.
☐ THE ISSUE FEE SET IN 37 C.F.R. 1.18 AT OR BEFORE MAILING OF THE NOTICE OF ALLOWANCE, PURSUANT TO 37 C.F.R. 1.31(b).
☐ ANY FILING FEES UNDER 37 C.F.R. 1.16 FOR PRESENTATION OF EXTRA CLAIMS.

THE FILING FEE HAS BEEN CALCULATED AS SHOW BELOW:

	(COL. 1) NO. FILED			(COL. 2*) NO. EXTRA	SMALL ENTITY			LARGE ENTITY	
					RATE	FEE		RATE	FEE
BASIC FEE:						\$380.00	OR		\$760.00
TOTAL CLAIMS:	21	-	20	1	X \$9 =	\$	OR	X \$18 =	\$18.00
INDEP. CLAIMS:	7	-	3	4	X \$39 =	\$	OR	X \$78 =	\$312.00
MULTIPLE DEPENDENT CLAIMS					+ \$130 =		OR	+\$260 =	\$0.00
*IF THE DIFFERENCE IN COL. 2 IS LESS THAN ZERO, ENTER "O" IN COL. 2.					TOTAL:	\$			\$1090.00

☒ THE COMMISSIONER IS HEREBY AUTHORIZED TO SUBMIT ALL CORRESPONDENCE RELATING TO THIS CASE TO THE CORRESPONDENCE ADDRESS LISTED BELOW.

By: Timothy L. McCutcheon

Date: September 1, 1999

Timothy L. McCutcheon, Esq., for
Carol Talkington Verser, Ph.D.
Registration No. 37,459
Heska Corporation
1613 Prospect Parkway
Fort Collins, Colorado 80525
Telephone: (970) 493-7272
Facsimile: (970) 484-9505

CORRESPONDENCE ADDRESS:

**Carol Talkington Verser, Ph.D.
Heska Corporation
1613 Prospect Parkway
Fort Collins, CO 80525**

METHOD AND COMPOSITION TO PROTECT AN OBLIGATE CARNIVORE FROM A DISEASE OF ABNORMAL CARBOHYDRATE METABOLISM

FIELD OF THE INVENTION

The present invention relates to a method to protect obligate carnivores from a
5 disease of abnormal carbohydrate metabolism, such as diabetes or obesity. In particular,
such animals are protected by feeding them a nutritionally balanced diet that includes a
high protein content, moderate fat content, and low carbohydrate content.

BACKGROUND OF THE INVENTION

Cats and dogs are the most common companion animals kept by man. As they are
10 both members of the biological order Carnivora, there is a tendency to assume that these
two carnivores have similar nutritional requirements. However, there are important
differences in the metabolism and nutritional requirements of cats and dogs; see, for
example, Morris et al., 1989, in *Waltham Symposium 7, Nutrition of the Dog and Cat* ed.
Burger et al., Cambridge University Press, pp. 35-66.

15 The Feloidae (Felids, Hyaenids, and Viverrids) diverged from the other members
of the order Carnivora relatively early in their evolutionary development. In contrast to
the Canoidae (Canids, Ursids, Procyonids and Mustelids), all members of the Feloidae are
flesh-eaters, i.e. strict or obligate carnivores. A comparison of the nutritional
requirements of cats and dogs as representative members of the Feloidae and Canoidae
20 supports the thesis that specialization consistent with the evolutionary influence of a strict
carnivorous diet has occurred in cats; see, for example, MacDonald et al., 1984, *Ann.*
Rev. Nutr. 4, pp. 521-562. A strict carnivorous diet implies the intake of a high protein,
moderate fat, and very low carbohydrate diet, following the composition of prey animals.

However, omnivorous species, like dogs, are adapted to both plant and animal food sources. Plants, unlike animals, have high carbohydrate stores in the form of starches.

One adaptation to a strictly carnivorous diet are differences in carbohydrate metabolism. For example, while the feline liver contains hexokinase, the enzyme
 5 responsible for the first step in glucose metabolism, it contains no glucokinase, which is a hexokinase that exhibits a significantly higher activity for the specific phosphorylation of glucose. Therefore, cats and other obligate carnivores might not be expected to be well adapted for the ingestion of high carbohydrate meals; see, for example, Morris et al., 1989, *ibid*. Additionally, the release of insulin from a cat's pancreas (insulin causes the
 10 cellular uptake of glucose from the blood) is dissimilar to that observed in most other species, appearing to be less responsive to glucose as a stimulus; see, for example, Curry et al., 1982, *Comparative Biochemistry and Physiology 72A*, pp. 333-338.

Another adaptation to a strictly carnivorous diet relates to the utilization of protein (made up of amino acids) and fat for energy production. Production of glucose from
 15 amino acids and fats is called gluconeogenesis. In an omnivore, gluconeogenesis occurs primarily in starvation situations, when the animal needs glucose to fuel its metabolism but can obtain it only from its own muscle protein; see, for example, Zubay, 1993, *Biochemistry*, Addison-Wesley. However, in an obligate carnivore, such as the cat, gluconeogenesis appears to be active at all times in the liver, regardless of nutritional
 20 status. Since an obligate carnivore normally has very low intake of carbohydrate, and its carbon sources are primarily protein and fat, it would be expected that the liver would be adapted for use of proteins as its primary source of glucose, rather than carbohydrate.

Additionally, it appears that cats, unlike omnivorous species, have limited ability to regulate the catabolic enzymes of amino acid metabolism. Therefore, when cats are fed a low protein diet, a high obligatory nitrogen loss results. Inability to down-regulate breakdown of amino acids accounts for the observed need for a significantly higher protein intake for cats relative to dogs; see, for example, Rogers et al., 1980, in *Nutrition of the Dog and Cat*, ed. R.S. Anderson, Oxford-Permagon Press, pp. 145-156.

Furthermore, cats and other obligate carnivores require animal source foods to meet their requirements for certain nutrients. For example, in contrast to dogs, cats cannot convert carotene from plants to Vitamin A; cats cannot synthesize niacin from tryptophan; cats cannot synthesize arachidonic acid from linoleic acid; and cats cannot synthesize sufficient taurine from cysteine. All of these nutrients can be found in a carnivorous diet; see, for example MacDonald et al., 1984, *ibid*.

Taken together, these data suggest that obligate carnivores, such as cats, are adapted to the use of protein not only for normal structural development and repair, but also as the primary source of energy via the process of gluconeogenesis.

Despite these metabolic differences between a cat, an obligate carnivore, and a dog, a omnivore, cat food formulations have traditionally been very similar to, and frequently derived from, dog food formulations. It has been known that cats have a higher requirement for protein, so cat food formulations have been modified to include higher concentrations of protein compared to dog foods. However, there have been a number of widespread health problems in cats related to such cat food formulations, demonstrating that food developed for dogs is not optimal for cats. For example, cats

began developing heart problems relating to lack of taurine in the diet. Taurine is only obtained from animal source protein since vegetable source protein does not contain taurine. Cats also began developing urinary stones, related to too much magnesium in the diet, and hypokalemia, a condition caused by low dietary potassium; see, for example

5 MacDonald et al., 1984, *ibid.* and Morris et al., 1989, *ibid.* Pet food companies have responded by supplementing their formulations to correct these problems caused by deficiencies in their cat food formulations. However, these problems demonstrate that there are very basic differences in dietary requirements between the cat, an obligate carnivore, and the dog, an omnivore, and that these differences are not fully addressed in

10 current cat foods.

Commercial cat foods today contain significant, even very high, levels of dietary carbohydrate from corn, wheat and other cereal grains. Dry formulation cat foods, in general, contain higher levels of grain carbohydrates than do canned varieties.

There has been a recent increase in the incidence of feline obesity and feline

15 diabetes in domesticated cats. Twenty percent of adult pet cats are thought to be obese, and feline diabetes is thought to affect one cat out of every four hundred; Panciera et al, 1990, *JAVMA* 197, pp. 1504-1508. Adult-onset diabetes, the most common form in the cat, is almost always insulin dependent and extremely difficult to regulate, even in the face of conscientious care by clinician and owner. The current therapy for feline adult-

20 onset diabetes is administration of insulin. Additionally, for feline obesity, the current therapy is much like that for the treatment of human obesity: it consists of a diet of lower

caloric density. Since fat has a higher caloric density than carbohydrate, typically carbohydrate is substituted for fat in order to lower caloric density.

Diabetes mellitus may occur either as a primary disease process or as a secondary complication caused by the destruction of beta cells or insulin resistance due to another
5 disease. In human disease, diabetes is subdivided into type 1 diabetes (insulin dependent, characterized by destruction of the insulin-secreting beta cells of the pancreas) and type 2 diabetes (non-insulin dependent, characterized by insulin resistance). Although there is strong evidence that both types occur in cats, type 2 diabetes appears to be much more frequent, and, in contrast to humans with type 2 diabetes, most cats with type 2 diabetes
10 are insulin dependent; see, for example Lutz et al. 1995, *Diabetes Mellitus* 25, pp. 527-549.

In type 2 diabetes, insulin secretion and insulin resistance are reduced compared to normal animals. For example, in diabetic cats and humans show markedly reduced or absent insulin secretion during the first phase of insulin response after an increase in
15 glucose, and a markedly delayed and often exaggerated insulin secretion during the second phase of the response. Impaired glucose tolerance is relatively common in cats. Additionally, in cats, marked suppression of insulin secretion may occur within days of a cat showing persistent marked hyperglycemia of approximately 540 milligrams (mg) per deciliter (dL) is present; see, for example Lutz et al., 1995, *ibid*. This phenomenon is
20 called glucose toxicity. In humans, insulin resistance, i.e. a state in which higher insulin concentrations are required to achieve a given amount of glucose uptake and utilization,

is determined genetically; however, in cats a predisposition to insulin resistance has not been demonstrated; see, for example, Rand, 1997, *Aust. Vet. Practit.* 27, pp. 17-26.

Diabetes in cats is currently difficult to treat and control. Treatment consists of either oral hypoglycemic drugs or insulin therapy. Sulfonylureas are the most common
5 class of oral hypoglycemic drugs. They act by both increasing insulin secretion from beta cells and the sensitivity of peripheral tissues to insulin, therefore, they are only useful if some functional beta cells are present. Insulin therapy requires careful control and monitoring of blood glucose levels, a challenge to clinician and cat owner. Only 30 to 50% of cats can be managed with oral hypoglycemic drugs. Additionally, approximately
10 15% of cats with diabetes are transient diabetics, meaning that therapy may be discontinued after several months or weeks; see, for example, Rand, 1997, *Aust. Vet. Practit.* 27, pp. 68-78.

Current thought on perspectives for treating feline diabetes focuses on understanding the role of the hormone amylin and glucagon-like peptide-1. Amylin
15 receptor antagonists are being developed to treat human type 2 diabetics, and are thought to be potentially useful for cats. Glucagon peptide-1 is also thought potentially useful in treating diabetic cats, Lutz et al., 1995, *ibid.* There remains, however, a need for a better method to protect cats from diseases of abnormal carbohydrate metabolism, including a method to maintain the well-being of such animals.

20 SUMMARY OF THE INVENTION

The current invention relates to the surprising discovery that feeding a cat a diet including low carbohydrate content, high protein content, and moderate fat content

improves that cat's health, or well-being. Such a diet is very similar to the natural diet that obligate carnivores, such as cats, have evolved to eat. This invention addresses all obligate carnivores, including all species from the Feloidae family and obligatory carnivores from the Canoidae family. Since an obligate carnivore is not adapted to

5 carbohydrate in its diet, and furthermore has an inability to regulate its hepatic enzymes as omnivores do, without being bound by theory, the inventor believes that the intake of dietary carbohydrate at levels seen in commercial cat foods would actually be harmful to an animal, causing depletion and/or suppression of insulin, and ultimately causing those most sensitive to these effects to become clinically diabetic. The inventor also believes

10 that obesity in an obligate carnivore, such as the cat, is caused, at least in part, by a high carbohydrate intake. High intake of carbohydrate by an animal adapted to almost exclusive intake of fat and protein appears to cause abnormal regulation of the hormones and enzymes that signal the fed and unfed state to the body. Further, this consumption of carbohydrates with the concomitant increase in circulating insulin causes the energy of

15 the diet to be stored as fat.

The present invention includes a method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism. The method includes the step of feeding the carnivore a nutritionally balanced diet that includes a low carbohydrate content; a high protein content, preferably animal source protein; and a moderate fat content. In one

20 embodiment, such a diet maintains the health, or well-being, of an obligate carnivore that is in good health. A preferred nutritionally balanced diet comprises a protein content of from about 25% to about 60% on a dry matter basis, a fat content of from about 15% to

about 60% on a dry matter basis, and a carbohydrate content of not more than about 12% on a dry matter basis. Another preferred nutritionally balanced diet comprises a protein content of from about 30% to about 70% on a dry matter basis, a fat content of from about 10% to about 40% on a dry matter basis, and a carbohydrate content of not more than about 12% on a dry matter basis. Another preferred nutritionally balanced diet includes animal meat wherein the balance of nutrients in the animal meat approximates the balance of nutrients in a total animal carcass.

The present invention also includes a method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism that includes the following steps: (a) producing a nutritionally balanced diet comprising a protein content of from about 25% to about 60% on a dry matter basis, a fat content of from about 15% to about 60% on a dry matter basis, and a carbohydrate content of not more than about 12% on a dry matter basis; and (b) feeding such a diet to the obligate carnivore. The present invention also includes a method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism that includes the following steps: (a) producing a nutritionally balanced diet comprising a protein content of from about 30% to about 70% on a dry matter basis, a fat content of from about 10% to about 40% on a dry matter basis, and a carbohydrate content of not more than about 12% on a dry matter basis; and (b) feeding such a diet to the obligate carnivore.

The present invention also includes such nutritionally balanced diets and methods to produce such diets. A preferred nutritionally balanced diet includes a protein content of from about 25% to about 60% on a dry matter basis, a fat content of from about 15% to

about 60% on a dry matter basis, and a carbohydrate content of not more than about 6%
on a dry matter basis. Another preferred nutritionally balanced diet includes a protein
content of from about 30% to about 70% on a dry matter basis, a fat content of from
about 10% to about 40% on a dry matter basis, and a carbohydrate content of not more
5 than about 12% on a dry matter basis.

The present invention also includes a method to protect an obligate carnivore from
a disease of abnormal carbohydrate metabolism, comprising feeding the obligate
carnivore a nutritionally balanced food composition that comprises a percentage of
carbohydrate on a dry matter basis that is not more than the highest percentage of
10 carbohydrate on a dry matter basis that will protect the obligate carnivore from such a
disease. In one embodiment, this method also includes feeding any nutritionally balanced
food composition that demonstrates a similar protective effect as a food composition
comprising animal meat wherein the balance of nutrients in said animal meat
approximates the balance of nutrients in a total animal carcass.

15 The present invention also includes a method to maintain the health of a healthy
obligate carnivore, said method comprising feeding the carnivore a nutritionally balanced
food composition that comprises a percentage of carbohydrate on a dry matter basis that
is not more than the highest percentage of carbohydrate on a dry matter basis that will
maintain the health of the carnivore. In one embodiment, the method includes feeding the
20 carnivore a nutritionally balanced food composition that demonstrates the same protective
effect as a food composition comprising animal meat wherein the balance of nutrients in
said animal meat approximates the balance of nutrients in a total animal carcass.

DETAILED DESCRIPTION OF THE INVENTION

The present invention includes a method to protect obligate carnivores, such as cats, from a disease of abnormal carbohydrate metabolism. This method includes the step of feeding the obligate carnivore a nutritionally balanced diet comprising a high protein, moderate fat, and low carbohydrate content. In one embodiment, the diet maintains the health, i.e., the well-being, of an obligate carnivore that is in good health, i.e., an animal that is not suffering from a disease of abnormal carbohydrate metabolism prior to being fed a diet of the present invention. Also included in the present invention are such nutritionally balanced diets.

As used herein, an obligate, or strict, carnivore is an animal that requires meat (i.e., animal flesh) in its diet. Such an animal can also eat other foods, but needs an animal source of protein, or supplements thereof, and preferably an animal source of fat, or supplements thereof, in order to maintain its health, i.e., well-being. Preferred obligate carnivores to feed in accordance with the present invention are Feloidae (e.g., Felids, Hyaenids, and Viverrids), as well as those Canoidae that are obligate carnivores (e.g., obligately carnivorous Mustelids). Preferred genera include, but are not limited to, *Felis*, *Panthera*, *Puma*, *Crocota*, *Hyaena* and *Putorius*, with the following species being even more preferred: *Felis domesticus*, *Felis silvestris*, *Panthera leo*, *Panthera tigris*, *Panthera pardus*, *Puma concolor*, *Crocota crocuta*, *Hyaena hyaena* and *Putorius furo*. It is to be noted that the terms Feloidae and cats are used interchangeably herein.

In one embodiment, the present invention relates to a treatment for feline adult-onset diabetes mellitus, a disease affecting as many as one out of every four hundred cats,

by feeding a nutritionally balanced diet comprising a very low carbohydrate content, a high protein content and a moderate fat content. While not being bound by theory, the inventor believes that the intake of dietary carbohydrate at levels commonly seen in commercial cat foods can be harmful to the cat, for example by causing depletion and or suppression of insulin secretion, ultimately causing those animals most sensitive to these effects to become clinically diabetic. This idea is new and surprising, since the belief in the pet food formulation art is that the amount of carbohydrate present in the diet is not important, as long as sufficient amounts of vital nutrients, such as protein, fat, trace minerals, vitamins and amino acids are present. Most formulations for adult cats contain high amounts of carbohydrate; see Table 1.

Table 1. Protein, fat, and carbohydrate contents of selected commercial cat foods, on a dry matter basis.

	Brand	Name of diet	formulation	% protein	% fat	% carbohydrate	% fiber
5	Ralston-Purina	CAT CHOW	dry	37	13.2	40.3	1.7
	Ralston-Purina	KITTEN CHOW	dry	41	13.8	35.5	1.8
	Ralston-Purina	CAT CHOW MATURE	dry	36.8	9.7	40.1	1.7
	Ralston-Purina	O.N.E. CHICKEN AND RICE FORMULA FOR CATS	dry	34.4	15.5	41.5	1.4
	Ralston-Purina	O.N.E. CHICKEN AND RICE FORMULA FOR KITTENS	dry	39.1	18.4	37.2	1.4
10	Ralston-Purina	PRO PLAN CAT GROWTH FORMULA	dry	36.5	22.7	34.5	1.2
	Ralston-Purina	PRO PLAN CAT ADULT FORMULA	dry	34.2	16.6	38.3	1.4
	Ralston-Purina	PRO PLAN CAT LITE FORMULA	dry	34.8	9.4	40.1	2.5
	Ralston-Purina	PRO PLAN TURKEY AND BARLEY FORMULA FOR CATS	dry	35.8	18.9	34.8	2.4
15	Hill's	PRESCRIPTION DIET FELINE C/D	dry wet	34.5 43.4	16.3 21.8	42.7 25.1	0.79 3.2
	Hill's	PRESCRIPTION DIET FELINE D/D	wet	39	28.1	26.2	1.5
	Hill's	PRESCRIPTION DIET FELINE H/D	wet	43.4	26.7	23.2	0.3
	Hill's	PRESCRIPTION DIET FELINE R/D (for obese cats)	dry wet	37.7 36.2	8.4 7.7	31.5 20.9	16.7 29.8
	Hill's	PRESCRIPTION DIET FELINE W/D	dry wet	39.2 41.1	9.5 16.6	36.5 23.7	9.0 12.3
	Hill's	HEALTH BLEND KITTEN	dry wet	36.1 43.2	23.7 33.1	32.7 16.6	1.1 0.6
20	Hill's	HEALTH BLEND FELINE	dry wet	32.7 41.1	20.2 22.3	41.2 28.7	1.1 1.0
	Hill's	HEALTH BLEND FELINE GERIATRIC	dry wet	33.3 41.2	18.6 20.6	41.1 28.8	1.7 4.1
	Hill's	SCIENCE DIET FELINE GROWTH	dry wet	37.1 49	26.8 36.2	29.1 6.9	1.2 0.6
	Hill's	SCIENCE DIET FELINE MAINTENANCE	wet dry	45.2 33.8	25.2 23	20.1 40	2.5 0.9
	Hill's	SCIENCE DIET LIGHT FORMULA FELINE MAINTENANCE	dry wet	40.8 44.9	9 12	35.6 27.8	8.4 9.2

Until the present invention, diabetic cats have typically been treated with either oral
 25 hypoglycemic drugs or exogenous insulin, and it had been thought that food intake had
 little effect on blood glucose in diabetic cats; see, for example, Martin et al, 1997, In

Proceedings of 15th American College of Veterinary Internal Medicine, p 670. The idea of treating diabetes by modifying the diet to remove most of the carbohydrate is new, and finding that blood glucose levels in diabetic cats are lowered when treated in accordance with the present invention is unexpected.

5 Another embodiment of the present invention relates to the treatment of feline obesity, an epidemic affecting as many as 20% of domestic cats, by feeding obese cats a nutritionally balanced diet comprising a very low carbohydrate, high protein and moderate fat content. Until the present invention, obesity in cats has been treated in a similar manner to treating obesity in humans, by, for example, lowering the caloric
10 density of the diet by increasing the amount of carbohydrate in the diet and lowering fat content. However, the cat's satiety response, as an obligate carnivore, is related to the amount of protein and fat the cat eats. Carbohydrate is not a natural dietary component in the cat's diet, so it would follow that carbohydrate will not provide the cat with a feeling of satiety. Thus, the cat is hungry and will overeat, causing weight gain. In accordance
15 with the present invention, a diet that contains moderate fat, with high animal protein content and low carbohydrate content, will be more effective at triggering the cat's satiety response, allowing it to maintain a more normal weight. The lower carbohydrate level of this diet will also result in lower levels of circulating insulin and less tendency to store dietary energy as fat.

20 It is to be noted that low carbohydrate diets currently exist in canned formulations only, and they are designed and positioned exclusively for the short term feeding of kittens. No such diet is claimed to be efficacious for the management and cure of feline

diabetes mellitus, obesity, or other diseases of abnormal carbohydrate metabolism.

Rather, the diets positioned as "diabetes diets" for cats have a high-fiber, moderate protein, moderate carbohydrate, and low fat profile that mimics the diets used to manage diabetes mellitus in dogs. In fact, no definitive research into the unique characteristics of feline diabetes and the resulting dietary implications has been done. Rather, what is believed and known about management of canine diabetes has been applied to cats, despite vast differences in the metabolism of obligate carnivores, such as cats, and omnivores, such as dogs. Thus, the invention of using a low carbohydrate diet with high protein and moderate fat to manage, cure, and prevent a disease of abnormal carbohydrate metabolism is different and new.

One embodiment of the present invention is a method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism by feeding the obligate carnivore a nutritionally balanced diet that includes a low carbohydrate, high protein and moderate fat content. As used herein, to protect from a disease means to prevent, control, cure, ameliorate or reduce the severity of said disease. In accordance with the present invention, to protect also includes to maintain the health, i.e., well-being, of an animal. It is to be noted that the term "a" or "an" entity refers to one or more of that entity; the terms "comprising", "including", and "having" can be used interchangeably. It should also be noted that the phrase "selected from the group consisting of" means any one or any combination of the members of the group.

Feeding includes food given orally as well as nutrition delivered by other routes, including, but not limited to, intravenous administration.

Nutrient on a dry matter basis is defined as the nutrient amount in a food composition after essentially all moisture is removed. Converting percent nutrients in a food composition to percent dry matter content is used to make direct comparisons of nutrient profiles of products with differing moisture contents. The term nutrient includes
 5 any ingredient that has a nutritive value for an animal, and includes the protein component, the fat component, and the carbohydrate component of a pet food as well as any other component(s) that is (are) necessary and/or is (are) thought to be necessary for the animal's health.

The term nutritionally balanced diet as used herein means a composition that,
 10 when fed as the sole source of food to an animal, contains sufficient kinds and amounts of nutrients to maintain an animal's health, i.e., well-being, over the course of the animal's life. A nutritionally balanced diet of the present invention includes a high protein content, a moderate fat content and a low carbohydrate content, each of which is described in more detail below. Such a nutritionally balanced diet also includes sufficient
 15 quantities of other ingredients, such as vitamins, minerals, amino acids, salts, and the like to maintain an animal's good health; appropriate ingredients and sufficient amounts to include in a nutritionally balanced diet are known to those skilled in the art. Examples of such ingredients include, but are not limited to, sodium chloride, potassium chloride, potassium citrate, calcium carbonate, iodized salt, choline chloride, taurine, iron oxide,
 20 zinc oxide, ferrous sulfate, copper sulfate, manganous oxide, sodium selenite, calcium iodate, vitamin E supplement, thiamine, niacin, calcium pantothenate, pyridoxine hydrochloride, riboflavin, folic acid, biotin and vitamin B12 supplement. Other

ingredients may be added in this composition as desired; such additives include flavoring agents, coloring agents, inorganic compounds, and fillers. A filler, or bulking agent, is defined as an ingredient that by addition to the composition, brings the composition to a one hundred percent composition. A filler or bulking agent comprises a non-digestible component, examples of which include, but are not limited to, non-digestible proteins, non-digestible fats, or non-digestible carbohydrates. Non-digestible carbohydrate is also known as fiber, examples of which include, but are not limited to, cellulose, peanut hulls, and soy fiber.

A disease of abnormal carbohydrate metabolism is a disease brought about by a concentration of carbohydrate in the blood which is sufficient to cause disease in an obligate carnivore. That is, the concentration of carbohydrate in the blood, typically measured as the blood glucose concentration, or blood glucose level, is abnormally high, thereby leading to disease. A blood glucose concentration of from about 90 mg glucose per dL blood to about 110 mg glucose per dL blood is considered normal in a cat. As such, blood glucose concentrations of greater than about 150 mg glucose per dL blood are considered indicators of an animal either having a disease of abnormal carbohydrate metabolism or being susceptible to such a disease. Since an obligate carnivore has evolved to exist on a diet composed of very low carbohydrate, the amount of carbohydrate the animal can tolerate before an abnormality, manifested as a disease, is lower than what is tolerated by an animal that is not adapted to exist on a very low carbohydrate diet. Examples of diseases of abnormal carbohydrate metabolism include, but are not limited to, diabetes and obesity.

The amount of carbohydrate present in a nutritionally balanced diet of the present invention is of key importance and should be sufficiently low so as not to cause, or lead to, a disease of abnormal carbohydrate metabolism in the fed animal. In one embodiment, the amount of carbohydrate in an obligate carnivore's diet, and preferably in a cat's diet, on a dry matter basis, preferably is less than about 20% carbohydrate. More preferred is a concentration less than about 15% carbohydrate, even more preferred is a concentration of less than about 12% carbohydrate, even more preferred is a concentration of less than about 10% carbohydrate, even more preferred is a concentration of less than about 8% carbohydrate, even more preferred is a concentration of less than about 6% carbohydrate, even more preferred is a concentration of less than about 5% carbohydrate, even more preferred is a concentration of less than about 4% carbohydrate, and even more preferred is a concentration of about 2% or less carbohydrate. In one embodiment, no carbohydrate is added to the cat's diet.

Carbohydrates can be understood to be monosaccharides, oligosaccharides, and/or high molecular weight polysaccharides. Monosaccharides are defined as any of a class of aldehyde or ketone derivatives of polyhydric alcohols, particularly of the pentahydric and hexahydric alcohols. Common monosaccharides include glucose, fructose, galactose, and mannose. Most carbohydrates in nature exist as polysaccharides, which are high molecular weight oligomers of monosaccharides linked by glycosidic bonds. Ingested polysaccharides are broken down into their component monosaccharides by enzymes that are specific for that particular polysaccharide. Carbohydrate may be supplied in the form

of vegetable starches, which can include but are not limited to cereal grains such as wheat, corn, barley and rice.

The protein components used to prepare the nutritionally balanced diet with high protein content can be supplied by ingredients such as animal source protein, including
5 but not limited to meat and meat by-products; plant protein sources supplemented with essential amino acids, such as taurine, and/or other protein-related nutrients, as needed, in an amount to be nutritionally balanced; or a combination thereof. By meat is meant the flesh of an animal, such as, but not limited to, cattle, swine, sheep, goats, horses, other hoofed-animals, other mammals, poultry, fish, and arthropods. Meat by-products
10 include, but are not limited to, lungs, kidneys, brain, livers, and stomachs and intestines freed of their contents. Preferred is a nutritionally balanced diet in which the protein ingredients are derived at least in part from animal source protein; more preferred is a nutritionally balanced diet in which the protein ingredients are derived completely from animal source protein. A preferred nutritionally balanced diet of the present invention is
15 one in which the protein content, on a dry matter basis, ranges from about 25% to about 70% protein, more preferably from about 30% to about 60% protein, more preferably from about 40% to about 55% protein, and even more preferably from about 48% to about 52% protein.

The fat in a nutritionally balanced diet of the present invention is one or more
20 triacylglycerols of fatty acids, and may be derived from an animal source of fat; a plant source of fat supplemented, as needed, with essential fatty acids or fat-associated or related nutrients, for example arachidonic acid, in an amount to be nutritionally balanced;

or a combination of animal source fat and vegetable source fat. Preferred is a nutritionally balanced diet wherein the fat ingredients are derived at least in part from animal source fat; and more preferred is a nutritionally balanced diet in which the fat ingredients are derived completely from animal source fat. A preferred nutritionally balanced diet of the present invention is one in which the fat content, on a dry matter basis, ranges from about 15% to about 60% fat, more preferably from about 25% to about 50% fat, more preferably from about 30% to about 45% fat, and even more preferably from about 35% to about 40% fat.

One aspect of the present invention is a method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism which comprises feeding the obligate carnivore a nutritionally balanced diet of animal meat. Such animal meat has a nutrient balance that approximates (i.e., is similar to) the balance of nutrients in a total animal carcass. An embodiment of this diet is a diet comprising whole ground rat carcasses. A whole ground rat carcass diet can be prepared by sacrificing adult rats, removing the gastrointestinal contents, and grinding the carcasses. This diet may be canned, refrigerated or frozen prior to feeding to the animal or fed fresh to the animal.

One embodiment of the present invention is an animal food composition comprising a nutritionally balanced diet. A preferred nutritionally balanced diet includes a protein content of from about 25% to about 60% on a dry matter basis, a fat content of from about 15% to about 60% of a dry matter basis, and a carbohydrate content of not more than about 12%, more preferably of not more than about 10%, more preferably of not more than about 8%, more preferably of not more than about 6%, more preferably of

not more than about 5%, more preferably of not more than about 4%, and even more preferably of not more than about 2% carbohydrate on a dry matter basis. Another preferred nutritionally balanced diet includes a protein content of from about 30% to about 70% on a dry matter basis, a fat content of from about 10% to about 40% of a dry matter basis, and a carbohydrate content of not more than about 12%, more preferably of not more than about 10%, more preferably of not more than about 8%, more preferably of not more than about 6%, more preferably of not more than about 5%, more preferably of not more than about 4%, and even more preferably of not more than about 2% carbohydrate on a dry matter basis. Another preferred nutritionally balanced diet is animal meat, such as a total animal carcass. Such compositions have utility in maintaining the health, or well-being, of an animal as well as for treating, curing, controlling or reducing disease in an animal as disclosed herein.

The present invention also includes a method to produce nutritionally balanced diets of the present invention. Such diets can be prepared as wet diets, which are typically canned, or dry diets, which are typically packaged in containers, such as bags. As used herein, a wet diet refers to any diet which is not a dry diet. Wet food diets of the present invention are produced using methods known to those skilled in the art. For example, wet food can be produced by mixing the desired ingredients with water sufficient for processing. In one embodiment, the mixture is typically heated in a suitable vessel to a temperature of from about 70°F to about 140° F. When heated to the appropriate temperature, the mixture is in the form of a thick liquid or paste. The thick liquid or paste is then placed into cans, lids are applied, and the cans are hermetically

sealed. The sealed can is then sterilized using any of a variety of methods known to those skilled in the art, such as by heating the can to about 230° F for an appropriate length of time.

Until the present invention, dry cat foods contained, on average, a significantly
5 higher amount of carbohydrate on a dry weight basis than wet cat food, and, indeed, the inventor is unaware of any commercial dry cat food formulation that contains a nutritionally balanced diet of the present invention. While not being bound by theory, it is believed that carbohydrate (e.g., starch) was added to facilitate the extrusion process by which currently commercial dry cat food is typically formulated after baking with steam
10 and pressure. There is, however, no reason to expect that it would be difficult to make a dry animal food composition comprising a nutritionally balanced diet of the present invention using methods known to those skilled in the art, including, but not limited to, drying (e.g., by baking) and extrusion. In one embodiment, a dry diet of the present invention is prepared by a process of slowly drying and baking a wet diet of the present
15 invention; for example, the wet diet preparation is placed in a heated oven, for example an oven at 225° F, for a sufficient amount of time to dry the wet preparation. The preparation is uncovered throughout this process, and is stirred as necessary during the drying process. After drying is complete, the mixture is cooled before being packaged. The resultant dry nutritionally balanced diet has a moisture content of about 10%,
20 compared to a starting moisture content of between 25% and 35%.

The present invention also includes a method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism that includes the step of feeding the

obligate carnivore a nutritionally balanced food composition that comprises a percentage of carbohydrate on a dry matter basis that is not more than the highest percentage of carbohydrate on a dry matter basis that will protect said obligate carnivore from said disease of abnormal carbohydrate metabolism. In one embodiment, the method includes
5 a food composition that has a similar protective effect as a food composition comprising animal meat wherein the balance of nutrients in said animal meat approximates the balance of nutrients in a total animal carcass.

The present invention also includes a method to maintain the health of a healthy obligate carnivore that includes the step of feeding the carnivore a nutritionally balanced
10 food composition that comprises a percentage of carbohydrate on a dry matter basis that is not more than the highest percentage of carbohydrate on a dry matter basis that will maintain the health of said healthy obligate carnivore. In one embodiment, the method includes a nutritionally balanced food composition that has a similar health maintenance effect as a food composition comprising animal meat wherein the balance of nutrients in
15 said animal meat approximates the balance of nutrients in a total animal carcass.

The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention.

EXAMPLES

Example 1

20 This Example demonstrates the ability of a nutritionally balanced diet of the present invention to treat a disease of abnormal carbohydrate metabolism in a cat.

A 9-week study was performed on an eight year old domestic shorthair cat with adult-onset, insulin dependent diabetes. This cat developed the clinical signs of diabetes mellitus at the age of five years. This cat, as an adult, had been fed commercial diets with high levels of carbohydrate; i.e., HILL'S SCIENCE DIET FELINE MAINTENANCE and HILL'S SCIENCE DIET FELINE LIGHT. This cat required treatment with insulin (ULTRA LENTE insulin SQ) to lower its blood glucose levels to a more normal range. However, as is common with cats with feline diabetes mellitus, this cat's disease was difficult to control. Frequent testing and careful titration of the cat's insulin dosage was required; even with these precautions, on two occasions, the cat lapsed into diabetic ketoacidosis, a complication caused by hyperglycemia.

At week 0 of the present study, the cat's diet was switched to a low carbohydrate, high meat protein, moderate fat diet. The diet, referred to herein as DIET 1, is a wet food formulation with 7% carbohydrate, 49% protein, and 36% fat on a dry matter basis. DIET 1, on a wet matter basis, has about 13% crude protein, 8% crude fat, 3.5 % ash, 1% crude fiber, 0.25% calcium, 0.2% phosphorus, 0.1% taurine, 0.04% magnesium and 72% moisture. At the time this diet began, this cat was being dosed with approximately 4 units of insulin every 12 hours. Dosage was adjusted over the course of this study in response to measured blood glucose level, with dosage being lowered by a half unit every day that the blood glucose was determined to be below 100 mg/dL. Table 2 shows the blood glucose levels and insulin dosage for the cat fed the low carbohydrate diet over a period of 66 days.

Table 2. Blood glucose levels and insulin dosage for treated cat.

DAY	DATE	Time	Glucose level	insulin dosage
0	5/25/1998	5:00 pm	230 mg/dL	3.5 units BID
4	5/29/1998	6:00 pm	200 mg/dL	3.5 units BID
9	6/3/1998	5:15 pm	180 mg/dL	3.0 units BID
11	6/5/1998	5:00 pm	90 mg/dL	2.5 units BID
13	6/7/1998	5:30 pm	70 mg/dL	2.0 units BID
15	6/9/1998	5:30 pm	30 mg/dL	1.0 unit BID
20	6/14/1998	6:00 pm	100 mg/dL	0.5 unit BID
27	6/21/1998	5:00 pm	68 mg/dL	0.0 unit BID
29	6/23/1998	6:00 pm	110 mg/dL	0.0 unit BID
36	6/30/1998	6:00 pm	324 mg/dL	1.0 unit BID
38	7/2/1998	not record.	30 mg/dL	0.0 unit BID
54	7/18/1998	12:00 pm	103 mg/dL	0.0 unit BID
60	7/24/1998	11:00 am	105 mg/dL	0.0 unit BID
66	7/30/1998	3:30 pm	97 mg/dL	0.0 unit BID

Results show the efficacy of the low carbohydrate, high protein and moderate fat diet for the treatment of feline diabetes mellitus in this cat. Hyperglycemia (measured by blood glucose levels) was reduced to approximately normal levels, and this level was maintained without exogenous insulin after an approximate four week treatment with the diet.

Example 2

This Example demonstrates the ability of a nutritionally balanced diet of the present invention to treat a disease of abnormal carbohydrate metabolism in several cats.

A study was set up to evaluate the effect of feeding nine cats with well characterized adult-onset, insulin-dependent diabetes mellitus of at least three month's duration a nutritionally balanced diet of the present invention. Cats enrolled into the study could be of any age, any breed or mixture of breeds, either gender, and of any

clinical severity except they could not be suffering from active ketoacidosis. At week 0 of the study, the cats began to be fed a nutritionally balanced diet of the present invention consisting of the canned formulation DIET 1, described in Example 1. No additional supplements, treats, table scraps, vitamins, minerals or any other nutritive substances
5 were fed during the study. The results, to date, of the study are as follows, reported on a cat-by-cat basis:

Case 1: 12 year old spayed female DSH (domestic short hair) cat. Overweight at 16 pounds. Diagnosed with diabetes mellitus two years ago. Started insulin injections 18 months ago. Prior to the start of the study, this cat was receiving 5 IU NPH insulin twice
10 daily and was eating HILL'S PRESCRIPTION DIET FELINE W/D. At the beginning of the study, the cat was taken off insulin; after two weeks on DIET 1, the cat exhibited stable blood glucose levels between 150 and 250 mg glucose per dL blood and showed clinical improvement.

Case 2: 16 year old male castrate, non-obese DSH cat. Diagnosed with diabetes
15 mellitus one year ago. On 2 IU NPH insulin twice daily prior to entering study and was eating HILL'S SCIENCE DIET FELINE MAINTENANCE. After a 3 week treatment with DIET 1, this cat was maintaining normal blood glucose levels without exogenously administered insulin.

Case 3: 9 year old male castrate, non-obese, domestic long hair, cat. Diagnosed
20 with diabetes mellitus one year ago. Prior to the start of the study, this cat was treated twice daily with 3 IU NPH insulin and was eating Ralston Purina's CAT CHOW. After

two weeks on DIET 1, this cat maintained normal blood glucose levels with a reduced treatment of 1 IU NPH insulin.

Case 4: 10 year old male castrate, non-obese, Siamese cat. Diagnosed with diabetes mellitus 6 months ago. Prior to the start of the study, this cat was treated twice daily with 8 IU NPH insulin and was eating HILL'S PRESCRIPTION DIET FELINE W/D. When DIET 1 was started, all insulin treatment was stopped; and after two weeks on the study diet, this cat was maintaining a normal blood glucose level.

Case 5: 14 year old male castrate DSH cat. Diagnosed with diabetes mellitus two months ago. Prior to the start of the study, this cat was treated twice daily with 5 IU NPH insulin and was eating HILL'S SCIENCE DIET FELINE MAINTENANCE. After two weeks on DIET 1, this cat was maintaining normal blood glucose levels without exogenous insulin.

Case 6: 13 year old male castrate Siamese cat. Moderately overweight at 13 pounds. Diagnosed with diabetes mellitus 6 months ago. Prior to the start of the study, this cat was receiving 8 IU NPH insulin twice daily and was eating HILL'S PRESCRIPTION DIET FELINE W/D. After 3 weeks on DIET 1, this cat was maintaining a normal blood glucose level with a reduced treatment of 4 IU NPH insulin twice daily.

Case 7: 14 year old male castrate DSH cat. Diagnosed with diabetes mellitus 3 years ago. Prior to the start of the study, this cat was treated twice daily with 6 IU NPH insulin and was eating HILL'S PRESCRIPTION DIET FELINE W/D. After one week

on DIET 1, this cat was maintaining a normal blood glucose level with a reduced treatment of 1 IU NPH insulin twice daily.

Case 8: Approximately 10-12 year old male castrate domestic long hair cat. Diagnosed with diabetes mellitus 2 years ago. On varying dosages of NPH insulin, ranging from 2-6 IU twice daily and was eating a variety of commercial grocery store cat foods. After one week on DIET 1, this cat was maintaining a normal blood glucose level with no exogenously administered insulin.

Case 9: 13 year old male castrate, slightly overweight, Balinese cat. Diagnosed with diabetes mellitus 5 months ago. Prior to start of study, this cat was receiving 6 IU ULTRA LENTE insulin twice daily and was eating Waltham, CNM UR, and Nestle's FRISKIES. After 3 weeks on DIET 1, this cat still required the same amount of insulin (6 IU) to maintain a normal blood glucose level, although this cat has shown better control and lower blood glucose levels than prior to the beginning of the study.

Case 10: 14 year old spayed female, DSH cat. Underweight at 6.6 pounds. Diagnosed with diabetes mellitus 3 years ago. Prior to the start of the study, this cat was receiving 6 IU NPH insulin twice daily and was eating HILL'S PRESCRIPTION DIET FELINE W/D. After 3 weeks on DIET 1, this cat was maintaining approximately normal blood glucose levels on a reduced amount of insulin, 1 IU, twice daily.

Case 11: 16 year old male castrate, non-obese, DSH cat. Diagnosed with diabetes mellitus 2 years ago. Prior to the start of the study, this cat was receiving 5 IU ULTRA LENTE insulin twice daily and was eating HILL'S PRESCRIPTION DIET FELINE

W/D. After 3 weeks on DIET 1, this cat was maintaining an approximately normal blood glucose level with a 0.5-0.8 IU ULTRA LENTE insulin twice daily.

Case 12: 8 year old male castrate DSH cat. Moderately overweight at 14 pounds. Diagnosed with diabetes mellitus 2 months ago. This cat was not receiving exogenous
 5 insulin injections, and had a blood glucose level in the 400 mg/dL range, and was eating IAMS cat foods and FRISKIES. After 1 week on DIET 1, this cat's blood glucose level dropped to approximately 300 mg/dL.

During the study, all cats ate the study diet well but did not show a tendency to overeat or to gain weight, except for the cat in case 10 who gained 0.5 pounds. All cats
 10 showed clinical improvement. These results indicate that a nutritionally balanced diet including low carbohydrate, high protein, moderate fat of the present invention improved the condition of cats suffering from diabetes.

Example 3

This example demonstrates the ability to produce a dry food formulation of a
 15 nutritionally balanced diet of the present invention.

A dry nutritionally balanced diet, defined as 4.77% carbohydrate, 32.4% fat, and 44.98% protein with the remainder of the composition being 5.51% ash, 10.94% moisture, and 1.4% fiber, was prepared as follows. A sample of DIET 1 was slowly dried and baked such that a starting moisture content of 25% to 35% was reduced to a moisture
 20 content of about 10%. The moist composition was spread on a tray, and placed in a 225° F convection oven for three to three and one half hours, uncovered. The mixture was

stirred once an hour during the drying process. After drying was complete, the mixture was cooled slowly for approximately one hour before being packaged.

Example 4

This example demonstrates the ability of a nutritionally balanced diet of the present invention to treat a disease of abnormal carbohydrate metabolism in several diabetic cats. This study was continued over several months.

A baseline glucose curve was taken and the cats were then switched to DIET 1, as described in Example 1. Glucose curves were performed in the following manner: each cat is fasted, and has no insulin injections for 12 hours. At time 0, the cat is given insulin and fed. A blood sample is taken every two hours following for ten hours, and the glucose amount is determined at each time point. In a normal cat, the glucose level of approximately 100 mg/dl will normally rise slightly upon food consumption, then drop slightly below 100 in response to the insulin released in response to the blood sugar rise. The blood glucose level should then return to normal.

15 Cat 1. "Sol" Akers

hour	baseline 3/1/98	Start diet 8/7/98	8/24/98	9/10/98	10/5/98	10/20/98	1/9/99	3/26/99
8:00 am	460	192	578	111	318	50	443	270
10:00 am	447	73	431	58	402	40	398	410
12:00 am	487	69	410	61	311	36	333	449
2:00 pm	418	76	399	70	262	35	319	427
4:00 pm	437	196	320	96	251	50	300	494
6:00 pm	445	160	334	100	230	N/D	N/D	N/D
insulin	8 units BID	4 units BID	3 units BID	4 units BID	3 units BID	5 units BID	3 units BID	11 units BID

This cat was taken off the study diet on February 16, 1999 due to recurrent pancreatitis. Before starting the study diet, this cat required 8 units of exogenously

administered insulin to maintain blood glucose under 500 mg/dl. However, for the first few months after the initiation of the study diet, this cat's glucose level was maintained at lower levels, allowing for a reduction of administered insulin. However, this cat lost its ability to maintain a stable blood glucose level, most probably due to pancreatitis.

5 Cat 2. "Scruffy" Hudson

10

hour	Baseline 7/1/98	Start diet 8/14/98	9/1/98	9/30/98	10/15/98	11/5/98	1/21/99	5/13/99
8:00 am	219	47	84	86	117	110	107	93
10:00 am	342	88	107	96	137	91	96	100
12:00 am	260	183	89	74	89	90	73	102
2:00 pm	109	213	101	68	79	81	66	106
4:00 pm	116	33	80	73	84	95	98	104
6:00 pm	150	160	100	N/D	N/D	N/D	N/D	N/D
insulin	6 units BID	1 unit BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID

Before starting the study diet, this cat required 6 units of exogenously

15 administered insulin to maintain a blood glucose level of 109-342 mg/dl. Upon starting the study diet, this cat was maintaining normal blood glucose levels with no exogenously administered insulin after an approximately 2-week treatment with the study diet. The cat has maintained normal blood glucose levels for approximately 9 months while being fed the study diet.

20 Cat 3. "Ollie" Moozakis

25

hour	Baseline 7/1/98	Start diet 8/11/98	8/24/98	9/9/98	9/23/98	10/5/98	11/2/98	1/28/99
8:00 am	160	80	99	82	84	87	84	93
10:00 am	130	28	63	79	100	82	100	83
12:00 am	150	63	78	85	78	84	88	79
2:00 pm	165	79	82	85	80	96	78	86
4:00 pm	180	75	83	87	82	91	85	83
6:00 pm	155	90	N/D	N/D	N/D	N/D	N/D	N/D
insulin	2 units BID	1 unit BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID

This cat was removed from the study due to poor dietary compliance. Before starting the study diet, this cat was able to maintain a blood glucose level of 130-180 with 2 units of exogenously administered insulin. Upon starting the diet, until removal from the study, the cat maintained normal blood glucose levels without exogenously administered insulin.

Cat 4. "Mabby" Couturier

hour	Baseline 8/14/98	Start diet 8/15/98	8/28/98	9/11/98	9/25/98	10/16/98	10/23/98	4/23/99
8:00 am	232	170	296	233	268	320	162	144
10:00 am	184	150	280	225	342	301	96	111
12:00 am	80	147	249	253	292	278	82	120
2:00 pm	60	126	216	215	202	270	105	132
4:00 pm	136	155	195	222	223	283	131	160
6:00 pm	146	160	213	248	248	273	136	N/D
insulin	5 units BID	3 units BID	0 units BID	0 units BID	0 units BID	0 units BID	1 unit BID	1 unit BID

This cat required 5 units of insulin to maintain blood glucose under 250 mg/dl. Upon starting the study diet, this cat maintained a stable blood glucose level between 200-300 mg/dl with no exogenously administered insulin. With administration of 1 unit of insulin, this cat was able to maintain a blood glucose level in the normal range. This result represents a reduction from 5 units of administered insulin, which represents a considerable improvement in the cat's ability to regulate its blood glucose levels.

Cat 5. "Leo" Yates

hour	Baseline 8/15/98	Start diet 8/24/98	9/10/98	9/24/98	10/8/98	10/22/98	11/5/98	1/16/99	4/19/99
8:00 am	381	154	99	117	111	110	116	168.8	123
10:00 am	301	39	104	131	103	85	122	125	128
12:00 am	200	55	98	92	106	94	95	102	91
2:00 pm	282	55	94	90	111	85	93	79	94
4:00 pm	300	92	95	88	128	80	85	85	116
6:00 pm	N/D	94	N/D	N/D	N/D	N/D	N/D	N/D	N/D
insulin	4 units BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID

10 This cat required 4 units of insulin before the start of the study diet. Even with 4 units of exogenously administered insulin, blood glucose levels were high, in the 200-400 mg/dl range. Upon starting of the study diet, this cat was maintaining normal blood glucose levels without exogenously administered insulin.

Cat 6. "Koshka" Carr

hour	Baseline 8/8/98	Start diet 8/9/98	8/22/98	9/9/98	10/3/98	10/24/98	1/15/99	3/22/99
8:00 am	486	426	497	500	442	370	455	310
10:00 am	425	250	448	382	418	308	459	209
12:00 am	375	220	369	388	371	236	354	182
2:00 pm	410	200	323	467	364	240	261	231
4:00 pm	417	300	356	459	455	242	219	268
6:00 pm	380	375	343	N/D	N/D	N/D	N/D	N/D
insulin	6 units BID	6 units BID	5 units BID	6 units BID	6 units BID	8 units BID	8 units BID	9 units BID

20 This cat required 6 units of insulin to maintain a blood glucose level under 500 mg/dl before the start of the study diet. Upon start of the study diet, the required insulin dosage was briefly lowered to 5 units; however, the cat was not able to maintain a lower blood glucose level and the administered insulin dosage was increased. The study diet did not have an apparent effect on the blood glucose levels of this cat.

Cat 7. "Frosty" Sharkoff

hour	Baseline 8/11/98	Start diet 8/12/98	8/29/98	9/13/98	9/27/98	10/10/98	10/31/98	1/23/99	3/6/99
8:00 am	584	380	35	101	71	76	89	74	79
10:00 am	432	300	21	71	69	82	93	73	72
12:00 am	231	265	51	N/D	74	79	N/D	78	68
2:00 pm	146	179	58	N/D	72	75	N/D	49	76
4:00 pm	126	120	70	N/D	N/D	N/D	N/D	N/D	N/D
6:00 pm	148	180	85	N/D	N/D	N/D	N/D	N/D	N/D
insulin	6 units BID	3 units A.M.	1 unit BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID	0 units BID

This cat required 6 units of insulin to maintain a blood glucose level below 584 mg/dl before feeding the study diet. Upon start of the study diet, this cat was maintaining normal blood glucose levels without exogenously administered insulin after a 4-week treatment with the study diet.

Cat 8. "Cassatt" Lambersor

hour	Baseline 8/3/98	Start diet 8/5/98	8/24/98	8/31/98	10/20/98
8:00 am	467	450	358	344	380
10:00 am	460	N/D	349	309	294
12:00 am	434	N/D	318	268	208
2:00 pm	406	20	322	325	237
4:00 pm	422	70	302	348	290
6:00 pm	420	80	304	356	310
insulin	6 units BID	6 units A.M.	0.3 units BID	0.8 units BID	1.5 units BID

This cat, before the start of the study diet, required 6 units of insulin to maintain a blood glucose level between 400 and 467 mg/dl. After two weeks on the study diet, this cat was maintaining stable blood glucose between 200 and 380 mg/dl with greatly reduced dosages of insulin, ranging from 0.3 units to 1.5 units.

Cat 9. "Boots" Allen

hour	Baseline 3/1/98	Start diet 8/15/98	8/31/98	9/13/98	9/27/98	10/19/98	11/1/98	1/10/99	4/26/99
8:00 am	335	184	72	105	99	76	87	103	97
10:00 am	189	147	81	102	113	87	83	84	96
12:00 am	108	27	95	72	87	90	75	73	101

2:00 pm	148	52	91	74	93	82	78	78	107
4:00 pm	112	63	2	81	103	69	82	82	89
6:00 pm	201	73	85	91	98	77	85	85	72
insulin	8 units BID	4 units A.M.	0 units	0 units	0 units	0 units	0 units	0 units	0 units

5 This cat required 8 units of exogenously administered insulin to maintain a blood glucose level of 112 to 335 mg/dl before the start of the study. After 2 weeks on the study diet, this cat was maintaining a normal blood glucose level with no exogenously administered insulin. This cat, while on the study diet, has maintained a normal blood glucose for 8 months.

10 Cat 10. "Bobby" Blakemore

hour	Baseline 8/20/98	Start diet 8/21/98	9/4/98	9/18/98	10/3/98	10/16/98	11/5/98	12/22/98	3/25/99
8:00 am	319	77	135	496	316	223	147	271	388
10:00 am	432	34	115	247	130	98	43	258	249
12:00 am	231	48	76	41	127	76	43	200	68
2:00 pm	146	33	107	43	227	97	126	199	90
4:00 pm	229	229	209	78	468	167	150	185	95
6:00 pm	320	320	250	110	N/D	N/D	N/D	N/D	N/D
insulin	4 units BID	2 units BID	1 unit BID	1 unit BID	1 unit BID	1 unit BID	1 unit BID	1 unit BID	1 unit BID

15 This cat, before the start of the study, required 4 units of exogenously administered insulin to maintain blood glucose of 146 to 432. After 2 weeks on the study diet, the cat was maintaining a lower, although somewhat unstable, blood glucose level with the reduced amount of exogenously administered insulin amount of 1 unit BID.

20

Cat 11. "Alley" Kadlub

hour	Baseline 8/15/98	Start diet 8/16/98	8/24/98	9/14/99	10/20/99
8:00 am	381	340	399	197	405
10:00 am	340	N/D	316	121	262
12:00 am	310	N/D	313	52	125
2:00 pm	280	30	293	125	134
4:00 pm	300	50	359	201	354
6:00 pm	320	65	373	272	376
insulin	4 units BID	4 units A.M.	0.5 units BID	1.2 units BID	1 unit BID

10 This cat before the start of the study, required 4 units of exogenously administered insulin to maintain a blood glucose level of 280-380 mg/dl. After 4 weeks on the study diet, the cat was maintaining a lower, if somewhat unstable, blood glucose level with the reduced amount of exogenously administered insulin amount of 1 unit BID.

15 In this Example, the study diet containing reduced carbohydrate (6.9% on a dry matter basis) was shown to be effective in reducing cat's dependence on insulin to regulate its blood glucose level. Out of 11 cats in the study, 4 of the 11 were able to maintain normal blood glucose levels without exogenously administered insulin, 4 of the 11 were able to maintain lower blood glucose levels with a reduced amount of exogenously administered insulin, 2 of 11 were dropped from the study (as noted), and
20 for one of the 11 cats, the study diet did not appear to have any effect on blood glucose levels or dependence on insulin.

Example 5

 This example demonstrates the ability of a nutritionally balanced diet of the present invention to treat a disease of abnormal carbohydrate metabolism in several
25 diabetic cats. The diet used in this Example is different from the diets used in the

previous Examples; in this Example, the diet contains a higher protein content and lower fat content than the diet used previously.

Cats were fed Hill's Prescription Diet W/D, referred to herein as the baseline diet, prior to starting on the study diet. A baseline glucose curve was taken while cats were being fed Hill's Prescription Diet W/D. Cats were then switched to the study diet, referred to herein as DIET 2, which was a wet canned food. The analysis of DIET 2 is as follows: 74% moisture; 16.3% protein; 6.42% fat (by acid hydrolysis); 0.7% crude fiber; 2.06% ash; and 0.52% carbohydrate. On a dry matter basis, by calculation, protein makes up 62.7% of the diet formulation; fat makes up 24.7% of the diet formulation; fiber makes up 2.7% of the diet formulation; ash makes up 7.9% of the formulation; and carbohydrate makes up 2% of the diet formulation.

Glucose curves were performed in the manner described in Example 4.

Cat 1. "Maggie" Radman. Started baseline diet on 2/7/99 and DIET 2 on 4/3/99.

Hours post start of curve	1/25/99	2/20/99	3/6/99	4/2/99	5/17/99	6/29/99
0	146	287	120	140	106	135
2	60	275	166	145	129	101
4	98	112	87	84	87	110
6	108	114	85	88	108	113
8	131	117	85	100	128	116
10	N/D	163	133	141	N/D	157
12	N/D	N/D	N/D	N/D	N/D	123
INSULIN AMOUNT	4 units BID	2 units BID	2 units BID	2 units BID	0 units	0 units
WT OF CAT (LB)	10.84	10.5	10.1	10	10	9.2

This cat, before the start of the study diet, required 4 units of insulin to maintain a normal blood glucose level. After initiation of the study diet, normal blood glucose levels were maintained without administration of exogenous insulin.

This cat lost 1.64 lbs while on the study diet, for a loss of 15% of its body weight.

- 5 Cat 2. "Sylvester" Hernandez. Started baseline diet on 2/7/99 and DIET 2 on 4/7/99.

Hours post start of curve	2/6/99	2/20/99	3/6/99	4/6/99	5/1/99	5/22/99	7/2/99
0	323	282	410	216	311	290	174
2	335	283	360	254	90	211	59
4	350	68	208	225	184	151	56
6	314	47	61	97	290	133	57
8	292	96	59	106	390	164	85
10	221	162	121	121	N/D	237	N/D
12	N/D	N/D	N/D	N/D	N/D	N/D	N/D
INSULIN AMOUNT	5 units BID	5 units BID	5 units BID	5 units BID	3 units BID	3 units BID	4 units BID
WT OF CAT (LB)	12.85	12.6	12.2	N/D	12.6	13.2	12.7

- 15 This cat, before the start of the study diet, required 5 units of insulin to maintain a
- 20 blood glucose level of 221 to 350 mg/dl. After initiation of the study diet, lower blood glucose levels (mostly within the normal range) were maintained with a lower dose (3 to 4 units) of exogenously administered insulin.

This cat lost minimal weight during the study; the loss amounted to only about 1.1% of its body weight.

Cat 3. "Bud" Moore. Started baseline diet on 2/7/99 and DIET 2 on 4/4/99.

	Hours post start of curve	2/6/99	2/20/99	3/6/99	4/2/99	4/4/99	5/15/99	6/28/99
5	0	392	302	253	369	154	298	315
	2	374	283	370	353	107	332	294
	4	246	237	342	338	85	338	287
	6	213	243	361	341	43	343	232
	8	252	249	360	258	99	364	253
	10	262	300	340	325	325	390	253
10	12	N/D	N/D	N/D	N/D	N/D	N/D	296
	INSULIN AMOUNT	5 units BID	5 units BID	?	3.5 units BID	2 units BID	2 units BID	1.75 units BID
	WT OF CAT (LB)	17.3	17.1	N/D	N/D	N/D	N/D	14.75

15 This cat, before the start of feeding the study diet, required 5 units of insulin to maintain a blood glucose level of 213 to 392 mg/dl. After initiation of the study diet, lower blood glucose levels (from 232 to 315 mg/dl) were maintained with a reduced insulin dosage of 1.75 units.

20 This cat lost weight during the study; at the last data point, the cat had lost 2.55 lbs, for a loss of 14.7% of its body weight.

Cat 4. "PJ" Womack. Started baseline diet on 2/7/99 and DIET 2 on 4/3/99

	Hours post start of curve	2/6/99	2/20/99	4/2/99	5/14/99	6/26/99
25	0	154	139	144	136	96
	2	102	146	109	165	92
	4	84	31	75	122	102
	6	74	55	87	101	81
	8	135	91	100	79	96
	10	179	115	103	96	N/D
	INSULIN AMOUNT	5 units BID	3 units BID	1 unit BID	0 units	0 units
30	WT OF CAT (LB)	16.8	16.5	15.25	14	13.8

This cat, before the start of study diet, required 5 units of insulin to maintain a blood glucose level in the normal range. After initiation of the study diet, this cat

exhibited the ability to maintain a normal blood glucose level without administration of exogenous insulin.

This cat lost weight during the study; at the last data point, the cat had lost 3 lbs, for a loss of 17.9% of its body weight.

5 Cat 5. "Sparky" Ovitt. Started baseline diet on 2/3/99 and DIET 2 on 3/31/99.

	Hours post start of curve	12/14/98	2/17/99	3/9/99	5/12/99	7/8/99
	0	99	16	24	18	26
	2	66	20	21	50	69
10	4	152	69	26	77	76
	6	192	99	68	84	86
	8	182	87	122	85	72
	10	180	108	96	70	75
15	INSULIN AMOUNT	2.5 NPH units BID	2.5 units BID	2.5 units BID	1 units BID	1 units BID
	WT OF CAT (LB)	12.8	12.4	12	11.8	11.5

This cat, before the start of the study diet, required 2.5 units of insulin to maintain a normal blood glucose level. After initiation of the study diet, this cat was able to maintain a normal blood glucose level at a reduced insulin dosage of 1 unit.

This cat lost weight during the study; at the last data point, the cat had lost 1.3 lb, for a loss of 10% of its body weight.

Cat 6. "Peter" Ovitt. Started baseline diet on 2/3/99 and DIET 2 on 3/31/99.

	Hours post start of curve	2/17/99	3/17/99	5/7/99	7/8/99
25	0	260	273	16	17
	2	240	288	N/D	24
	4	143	312	26	62
	6	121	301	70	78
	8	103	307	104	108
30	10	88	324	129	125
	INSULIN AMOUNT	2 units BID	2 units BID	5 units BID	4.5 units BID
	WT OF CAT (LB)	11.7	11.7	11.6	13.4

This cat, before the start of the study diet, required 2 units of insulin to maintain a blood glucose level ranging from 88 to 260 mg/dl. After initiation of the study diet, this cat's insulin level was increased to bring down the blood glucose into the normal range. An effect of the study diet on allowing the cat to maintain a lower blood glucose level with a lower dosage of insulin is not apparent in this cat.

This cat did not lose weight during the study; it had, by the last data point, gained 1.7 lbs, for a gain of 14.5% of its body weight.

Cat 7. "Sassie" Bailey. Started baseline diet on 2/3/99 and DIET 2 on 3/31/99.

Hours post start of curve	12/13/98	2/9/99	3/10/99	3/30/99	5/11/99	7/7/99
0	187	156	219	255	357	150
2	158	123	145	277	253	86
4	99	80	63	213	179	58
6	100	82	132	133	191	138
8	165	78	212	15	209	186
10	201	74	210	136	214	206
INSULIN AMOUNT	4 units BID	4 units BID	4 units BID	4 units BID	4 units BID	3 units BID
WT OF CAT (LB)	11.5	11.5	11.8	11.8	11.4	N/D

This cat, before the start of the study diet, required 4 units of insulin to maintain a normal blood glucose level. After initiation of study diet, this cat was able to maintain a normal blood glucose level with a reduced amount of insulin (3 units).

This cat maintained its weight during the study.

Cat 8. "Daisy" Stender. Started baseline diet on 2/2/99 and DIET 2 on 3/2/99.

	Hours post start of curve	1/29/99	2/18/99	3/2/99	3/9/99	3/23/99	4/7/99	5/4/99	5/25/99
5	0	87	40	36	65	88	68	66	73
	2	79	54	29	57	61	48	65	86
	4	56	72	26	46	41	89	74	78
	6	54	48	36	38	66	102	67	73
	8	70	55	37	61	42	74	65	71
10	10	69	69	54	58	42	N/D	74	67
	INSULIN AMOUNT	4 units BID	?	2 units BID	0	0	0	0	0
	WT OF CAT (LB)	17.4	N/D	N/D	N/D	N/D	N/D	N/D	15.2

- 15 This cat, before the start of the study diet, required 4 units of insulin to maintain a normal blood glucose level. After initiation of the study diet, this cat was able to maintain a normal blood glucose level without exogenously administered insulin.

This cat lost weight during the study; at the last data point, the cat had lost 2.2 lb, for a loss of 12.6% of its body weight.

- 20 Cat 9. "Bud" Price. Started baseline diet on 2/2/99 and DIET 2 on 3/2/99.

	Hours post start of curve	1/29/99	2/18/99	3/2/99	3/9/99	3/23/99	4/24/99	5/4/99	5/25/99
	0	143	233	85	137	329	209	168	171
	2	120	337	36	196	358	59	256	167
25	4	303	137	39	232	295	36	111	96
	6	313	45	25	155	260	14	87	81
	8	370	70	51	217	308	172	87	90
	10	388	86	50	211	274	N/D	111	95
30	INSULIN AMOUNT	6 units BID	5 units BID	4 units BID	0 units	0 units	2 units BID	1 units BID	1 unit BID
	WT OF CAT (LB)	17.8	N/D	N/D	N/D	N/D	N/D	13.7	13.5

This cat, before the start of the study diet, required 6 units of insulin to maintain a blood glucose level in the range of 120-388 mg/dl. After initiation of the study diet, this

cat was able to maintain a normal blood glucose level with a reduced level of administered insulin (1 unit).

This cat lost weight during the study; at the last data point, the cat had lost 4.3 lbs, for a loss of 24.1% of its body weight.

- 5 Cat 10. "Tom" McBride. Started baseline diet on 2/2/99 and DIET 2 on 3/2/99.

Hours post start of curve	1/29/99	2/18/99	3/2/99	3/23/99	4/13/99
0	212	377	256	346	356
2	229	275	48	218	210
4	161	229	36	185	170
6	47	256	21	238	235
8	69	301	54	245	294
10	187	303	181	289	N/D
INSULIN AMOUNT	4 units BID	4 units BID	3 units BID	4 units BID	4 units BID
WT OF CAT (LB)	14.6	N/D	14.25	N/D	N/D

- 10
- 15 This cat, before the start of the study diet, required 4 units of insulin to maintain a blood glucose level in the range of 47 to 230 mg/dl. After initiation of the study diet, this cat still required 4 units of insulin to maintain its blood glucose in approximately the same range. An effect of the study diet on the cat's ability to maintain its blood glucose with lower dosing of insulin is not apparent in this cat.

- 20 This cat's weight was not recorded at the last two data points. However, the data points captured indicate that this cat is maintaining its weight on the study diet.

Cat 11. "Sammy" Miller. Started baseline diet on 2/2/99 and DIET 2 on 3/2/99.

	Hours post start of curve	1/29/99	2/18/99	3/2/99	3/9/99	3/23/99	5/4/99	5/25/99
5	0	57	73	96	71	106	68	75
	2	139	60	74	80	85	83	69
	4	165	42	54	69	66	61	97
	6	149	23	51	54	42	76	78
	8	116	56	47	21	54	96	83
	10	117	68	57	49	83	75	76
10	12	131	N/D	N/D	N/D	N/D	N/D	N/D
	INSULIN AMOUNT	5 units BID	4 units BID	4 units BID	0 units BID	0	0	0
	WT OF CAT (LB)	14.9	N/D	N/D	15.4	14.8	14.9	14.2

15 This cat, before the start of the study diet, required 5 units of insulin to maintain a normal blood glucose level. After initiation of the study diet, this cat was able to maintain a normal blood glucose level without exogenously administered insulin.

This cat's weight stayed relatively constant throughout the study.

Cat 12. "Spooky" Coit. Started baseline diet on 2/2/99 and DIET 2 on 3/2/99.

20	Hours post start of curve	2/2/99	2/18/99	3/2/99	3/9/99	4/13/99	5/4/99	5/25/99
	0	89	110	220	209	306	283	58
	2	70	105	200	124	274	177	38
	4	175	74	72	66	192	48	81
25	6	354	180	35	16	144	40	67
	8	455	245	51	78	129	72	91
	10	453	210	94	81	190	62	N/D
	INSULIN AMOUNT	9 units BID	9 units BID	4 units BID	4 units BID	2 units BID	3 units BID	2 units BID
30	WT OF CAT (LB)	12.5	N/D	13.25	N/D	N/D	N/D	N/D

This cat, before the start of the study diet, required 9 units of insulin to maintain a blood glucose level that ranged from 70 to 453 mg/dl. After initiation of the study diet, by the last time point this cat was able to maintain a normal blood glucose level with a much reduced dosage of insulin (2 units).

This cat's weight was not recorded throughout the study, therefore no conclusions are drawn with regarding to weight.

Cat 13. "Punkin" Kunze. Started baseline diet on 2/1/99 and DIET 2 on 3/1/99.

	Hours post start of curve	12/21/98	2/15/99	3/1/99	4/12/99	5/14/99
5	0	306	116	208	321	275
	2	322	98	268	339	338
	4	268	126	258	332	335
	6	229	162	266	323	328
	8	278	183	276	299	240
10	10	N/D	206	257	307	306
	INSULIN AMOUNT	3 units BID	3 units BID	0	0	0
	WT OF CAT (LB)	10.7	11.46	12.24	11.04	10.12

This cat, before the start of the study diet, required 3 units of insulin to maintain its blood glucose level of 278 to 322 mg/dl. After initiation of the study diet, by the last time point this cat was able to maintain a blood glucose level in approximately the same range without administered insulin.

This cat's weight fluctuated throughout the study, but by the last time point was quite similar to the cat's weight at the beginning of the study.

Cat 14. "Santino" Fitzpatrick. Started baseline diet on 2/1/99 and DIET 2 on 3/1/99.

	Hours post start of curve	12/21/98	2/15/99	3/1/99	4/12/99	5/14/99	7/12/99
20	0	343	404	592	481	453	587
	2	271	277	454	249	278	371
	4	160	169	394	171	238	279
25	6	119	99	359	120	204	237
	8	204	57	274	142	161	183
	10	N/D	43	166	159	161	163
	12	N/D	N/D	112	N/D	187	116
30	INSULIN AMOUNT	8 units BID	10 units BID	11 units BID	8 units BID	9 units BID	10 units BID
	WT OF CAT (LB)	10.5	10.78	10.18	10.26	10.4	11.4

This cat, before the start of the study diet, required 8 units of insulin to maintain its blood glucose level below 350 mg/dl. After initiation of the study diet, this cat required 10 units of insulin to maintain its blood glucose in approximately the same range. An effect of the study diet on the cat's ability to maintain its blood glucose with lower dosing of insulin is not apparent in this cat.

This cat's weight fluctuated throughout the study, remaining relatively constant until the last time point, where this cat had gained 0.9 lb, a gain of 8.5% of its body weight.

Cat 15. "Boots" Carr. Started baseline diet on 2/1/99 and DIET 2 on 3/1/99.

Hours post start of curve	12/21/98	2/15/99	3/1/99	4/12/99	5/14/99
0	181	394	339	355	369
2	195	339	346	339	378
4	98	274	362	335	389
6	38	156	323	310	379
8	49	210	311	286	333
10	71	211	330	320	333
INSULIN AMOUNT	4 units BID	3 units BID	3 units BID	0	0
WT OF CAT (LB)	27	25.22	25.28	21.66	19.39

This cat, before the start of the study diet, required 4 units of insulin to maintain its blood glucose in a normal range. After initiation of the study diet, this cat exhibited stable blood glucose levels between 333 and 389 mg per dl without exogenously administered insulin.

This cat lost weight during the study; it lost a total of 7.6 lbs, for a loss of 28.2% of its body weight.

In summary, fifteen cats were included in the study and four cats (data not shown) were removed from the study. Of the fifteen cats fed the study diet, twelve showed

improvement, measured by reduction in the amount of insulin required to allow the cat to maintain a lowered blood glucose level. Three of the fifteen cats showed no improvement upon feeding of the study diet. Of the twelve cats that showed improvement, four of the cats were able to maintain normal blood glucose levels without administered insulin; two of the cats were able to maintain stable, although higher than normal blood glucose levels, without administered insulin; and six of the fifteen cats were able to maintain a normal blood glucose level with a reduced amount of administered insulin relative the amount the cat required before initiation of the study.

Seven of the fifteen cats included in the study lost weight while on the study diet. Five of the fifteen cats maintained their weight while on the study diet; and two cats gained weight on the study diet. One cat of the fifteen did not have enough recorded data points to be able to draw a conclusion about weight loss.

While various embodiments of the present invention have been described in detail, it is apparent that modifications and adaptations of those embodiments will occur to those skilled in the art. It is to be expressly understood, however, that such modifications and adaptations are within the scope of the present invention, as set forth in the following claims.

What is claimed is:

1. A method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism, said method comprising feeding said obligate carnivore a nutritionally balanced diet comprising a protein content of from about 30% to about 70%
5 on a dry matter basis, a fat content amount of from about 10% to about 40% on a dry matter basis, and a carbohydrate content of not more than about 12% on a dry matter basis.
2. The method of Claim 1, wherein said obligate carnivore is selected from the group consisting of Felids, Hyaenids, Viverrids, and obligately carnivorous
10 Mustelids.
3. The method of Claim 1, wherein said obligate carnivore is selected from the group consisting of *Felis domesticus*, *Felis silvestris*, *Panthera leo*, *Panthera tigris*, *Panthera pardus*, *Puma concolor*, *Crocota crocuta*, *Hyaena hyaena*, and *Putorius furo*.
4. The method of Claim 1, wherein said carbohydrate content is not more
15 than about 6% on a dry matter basis.
5. The method of Claim 1, wherein said carbohydrate content is not more than about 2% on a dry matter basis.
6. The method of Claim 1 wherein said disease is selected from the group consisting of obesity and diabetes mellitus.
- 20 7. The method of Claim 1, wherein said method maintains the health of an obligate carnivore in good health.

8. The method of Claim 1, wherein said nutritionally balanced diet is selected from the group consisting of a dry diet and a wet diet.

9. A method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism, said method comprising:

(a) producing a nutritionally balanced diet comprising a protein content of from about 30% to about 70% on a dry matter basis, a fat content amount of
5 from about 10% to about 40% on a dry matter basis, and a carbohydrate content of not more than about 12% on a dry matter basis; and

(b) feeding said diet to said obligate carnivore.

10. A method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism, said method comprising feeding a nutritionally balanced diet comprising animal meat wherein the balance of nutrients in said animal meat approximates the balance of nutrients in a total animal carcass.

11. A method to maintain the health of a healthy obligate carnivore, said method comprising feeding said obligate carnivore a nutritionally balanced diet selected from the group consisting of: a nutritionally balanced diet comprising a protein content of from about 30% to about 70% on a dry matter basis, a fat content amount of from
5 about 10% to about 40% on a dry matter basis, and a carbohydrate content of not more than about 12% on a dry matter basis; and a nutritionally balanced diet comprising animal meat wherein the balance of nutrients in said animal meat approximates the balance of nutrients in a total animal carcass.

12. An animal food composition for protecting an obligate carnivore from a disease of abnormal carbohydrate metabolism, said composition comprising a nutritionally balanced diet having a protein content of from about 30% to about 70% on a dry matter basis, a fat content amount of from about 10% to about 40% on a dry matter basis, and a carbohydrate content of not more than 6% on a dry matter basis.

13. The method of Claim 12, wherein said carbohydrate content is not more than 6% on a dry matter basis.

14. The composition of Claim 12, wherein said carbohydrate content is not more than about 4% on a dry matter basis.

15. The composition of Claim 12, wherein said carbohydrate content is not more than about 2% on a dry matter basis.

16. The composition of Claim 12, wherein said composition is selected from the group consisting of a dry composition and a wet composition.

17. The composition of Claim 12, wherein said composition maintains the health of a healthy obligate carnivore.

18. A method to protect an obligate carnivore from a disease of abnormal carbohydrate metabolism, said method comprising feeding said obligate carnivore a nutritionally balanced food composition that comprises a percentage of carbohydrate on a dry matter basis that is not more than the highest percentage of carbohydrate on a dry matter basis that will protect said obligate carnivore from said disease of abnormal carbohydrate metabolism.

19. The method of Claim 18, wherein said nutritionally balanced food composition has a similar protective effect as a food composition comprising animal meat wherein the balance of nutrients in said animal meat approximates the balance of nutrients in a total animal carcass.

20. A method to maintain the health of a healthy obligate carnivore, said method comprising feeding said obligate carnivore a nutritionally balanced food composition that contains a percentage of carbohydrate on a dry matter basis that is not more than the highest percentage of carbohydrate on a dry matter basis that will maintain
5 the health of said healthy obligate carnivore.

21. The method of Claim 20, wherein said nutritionally balanced food composition has a similar health maintenance effect as a food composition comprising animal meat wherein the balance of nutrients in said animal meat approximates the balance of nutrients in a total animal carcass.

